



Original Article

Lived Experiences of Elderly Patients with Coronary Artery Disease: A Hermeneutic Phenomenological Study

Majideh Heravi-Karimooi¹, Nahid Rejeh¹, Mohammad Abbasi^{2*}

¹ Elderly Care Research Center, School of Nursing and Midwifery, Shahed University, Tehran, Iran

² Department of Nursing, School of Nursing and Midwifery, Qom University of Medical Sciences, Qom, Iran

ABSTRACT

Article history

Received 5 March 2017

Accepted 13 May 2017

Introduction: Coronary artery disease is the most common type of cardiovascular disease. Despite the increasing number of patients with coronary artery disease, the lived experiences of these patients have remained unknown. The purpose of the study was to understand the lived experiences of elderly patients with coronary artery disease.

Methods: This qualitative research was conducted using an interpretive phenomenological approach. Participants included thirteen elderly patients (9 males and 4 females) with coronary artery disease selective of purposive sampling method from Coronary Care Unit of Shahid Beheshti Hospital in Qom, 2016. Data were collected through semi-structured interviews between 55 - 80 minutes. The interviews were recorded and transcribed. The Van Manen's proposed six steps was used to analyze the data.

Results: The main theme of study was "living with ups and downs". It included sub-themes of 1) losing calmness, 2) trying to achieve calmness through spirituality, 3) losing health, 4) trying to promote health through modification of lifestyle, 5) perceiving family support, 6) living in fear, 7) living with a damaged heart".

Conclusion: Findings of the research indicated that the coronary artery disease led patients to experience ups and downs for the participations. After the disease, they were trying to achieve calmness through some strategies. Members of the treatment team, especially nurses should provide self-care education for to make them be have successfully in this field.

Keywords: Experience, Coronary Artery Disease, Qualitative Study

Citation: Heravi-Karimooi M, Rejeh N, Abbasi M. Lived experiences of elderly patients with coronary artery disease: a hermeneutic phenomenological study. *Elderly Health Journal*. 2017; 3(1): 28-34.

Introduction

Cardiovascular diseases are the most prevalent serious disease and the most problem of health which threatens human life in the world (1). These diseases are the most common causes of death in the developed and developing countries and the numbers of suffering people from them are increasing (2). Based on the available reports, cardiovascular diseases are subjective to the most common cause of death in Iran (3). Coronary artery disease is the most common type of the cardiovascular disease which has made the following coronary artery atherosclerosis (1, 4).

Suffering from the coronary artery disease causes notable changes in all aspects of patients' lives. They

report several experiences after the onset of disease as well as different signs and symptoms of angina (5, 6). Elderly people experience fear, feeling of insecurity, as well as lack of confidence in the present and future after myocardial infarction (7).

Fear of death during sleep and fear of pain during walking have affected various aspects of their lives (3, 8, 9). Through a review on researches conducted in Iran, it was concluded that less quantitative researches have been conducted in Iran about experience of living with coronary artery disease. Furthermore, the conducted studies evaluated a few aspects of living

* **Corresponding Author:** Department of Nursing, School of Nursing and Midwifery, Qom University of Medical Sciences, Qom, Iran. **Tel:** +982537835588, **Email address:** mohamad_abbasi55@yahoo.com

with coronary artery diseases mostly rate of outbreak and risk factors for cardiovascular diseases (10).

Despite previous study in Iran (11) these experiences have not yet been examined deeply and enough cognition is not available in this field. Therefore, it seems necessary to conduct a qualitative research to explain experiences of elderly patients with coronary artery disease deeply.

Cognition of these patients' experiences leads to creation of a unique knowledge in this field and provides some information for members of the treatment team especially nurses about training, caring, and consulting this type of patients. The purpose of the study was to explore the lived experiences of elderly patients with coronary artery disease.

Methods

The qualitative study was performed using a hermeneutic phenomenological approach, a most commonly method for understanding meaning of lived experience. Van Manen believed that knowledge and understanding come through analyzing the reflective descriptions of people with relevant experience. Van Manen's proposed six steps were used to conduct this study (12). In Van Manen's opinion hermeneutic phenomenology, a combination of description and interpretation, expresses that description and interpretation are not separated from each other. It should be noted that Van Manen's six steps are as a guidance to conduct the research and there is no obligation to observe the order in the steps. Each activity is explained in the following.

The first step: Turning to the nature of lived experience

Van Manen expresses that the first step in studying a phenomenon is the researcher's interest in nature of that research. The researcher's attention was drawn to the phenomenon of the lived experiences of elderly patients with coronary artery disease since he was working in CCU of a hospital in Tehran.

The second step: Investigating experience as we live it

Patients who referred to Shahid Beheshti Hospital in Qom city, Iran in 2016, underwent a purposive selection of participants with maximum variation sampling; nine men and four women. The inclusion criteria were willingness to participate in the research, talk about their lived experiences and age 65 years or older; and people with severe spiritual-psychological problems and physical diseases were excluded. Demographic characteristics of the study participants are shown in Table 1.

Deep semi-structured face to face interviews were carried out individually to collect data conducted in 55 to 80 minutes sessions in coronary care unit and heart unit of Shahid Beheshti Hospital of Qom. Interviews started with a general question about cardiovascular disease. The researcher let the participants to say their real experiences about coronary artery disease through his silence. In the case of needing more perception and explanation, the researcher used follow-up questions such as "would you please explain more about the issue" or "what was your purpose about the part which you said". Interviews were continued until data saturation was achieved, in the study after eleven interviews, the interviews ended when no new data was resulted.

Table 1. Demographic characteristics of the study participants

Participant Number	Marital status	Occupation status	Age	Sex	Time after disease (y)
1	Married	Retired	66	Male	1
2	Married	Farmer	71	Male	5
3	Married	Housekeeper	62	Female	6
4	Married	Self-employed	64	Male	9
5	Married	Housekeeper	71	Female	8
6	Married	Housekeeper	65	Female	7
7	Married	Retired	69	Male	6
8	Married	Self-employed	66	Male	4
9	Married	Housekeeper	61	Female	3
10	Married	Self-employed	65	Male	6
11	Married	Self-employed	72	Male	7
12	Married	Farmer	71	Male	2
13	Married	Retired	73	Male	4

The third step: Reflecting on essential themes

Thinking in inherent themes was descriptor of the considered phenomenon. Researcher determined the main themes for the phenomenon of elderly patients' experiences with coronary artery disease through thematic analysis. Van Mannen has proposed three approaches for thematic analysis including holistic approach, selective approach, and detailed approach. It should be noted that holistic and selective approaches were applied in this study. In the process of data analysis, after converting audio files to texts, texts were studied several times accurately and general perception from participants' experiences was obtained through holistic approach, then researcher wrote some paragraphs and studied them several times accurately) 12). He selected those sentences and phrases that described experiences of elderly patients with coronary artery disease and extracted them as semantic units. This semantic units were then written in a separate paper. and placed in a separate category according to their semantic similarity, in order to obtain more general and abstract phrases.

The fourth step: Hermeneutic phenomenological writing

Van Mannen's approach is the art of writing and rewriting. In this stage the purpose was achieving a strong description of the phenomenon of elderly patients' experiences with coronary artery disease.

The fifth step: Maintaining a strong and oriented nursing relation to the Phenomenon

This stage is for maintenance of a strong and directional communication with the considered phenomenon. In this stage, in order to avoid getting away from the purpose of study, the researcher considered the main research question to maintain the strong and directional communication with the phenomenon.

The sixth step: Balancing the research context by considering parts and whole

Van Mannen is making a balance in the field of linking whole and detail. He worked on this stage by continues review of whole and detail through holistic and selective approaches and according to the main research question.

According to Lincoln and Guba, the authors verified the accuracy of the data with the four criteria of validity, reliability, credibility, and transferability. An effective communication was made with the participants based on trust to guarantee the validity of study (12). In order to increase the reliability of the data and create a deeper understanding of the phenomenon, the authors used prolonged exposure to the subject and obtained data, reviewing the data and matching them with the experiences of the participants and the observers' comments. To confirm the data, texts of the interviews were given to patients after analysis. Also, the research process was evaluated step by step through participation of the research team as well as other experts.

Ethical considerations

This article is resulted from the approved project of Shahed University with ethical code of IR.Shahed.REC 1395.108. The researchers received the allowance to conduct the study from the related authorities by introducing themselves to Shahid Beheshti Hospital in Qom city. Then, they selected the participants and received verbal informed consent from them. The researchers committed to keep their personal information, interviews, and anonymity confidential.

Results

The research was conducted in order to perceive, identify, and interpret the lived experiences of elderly patients with coronary artery disease. Findings of the research indicated that the main theme of lived experiences of elderly patients with coronary artery disease was "living with ups and downs" including sub-themes of 1) losing calmness 2), trying to achieve calmness through spirituality, 3) losing health, 4) trying to promote health through modification of lifestyle, 5) perceiving family support, 6) living in fear, and 7) living with a damaged heart.

1. Losing Calmness

Most of the participants expressed that they had lost their calmness after disclosure of their cardiovascular disease and also hospital discharge. In their expression, they had suffered from worry, and anxiety, thought seriously about importance of heart in life and the fact that if the heart has problem, their lives might be in danger. They had lost their calmness by reviewing their heart task and further probabilities such as cardiac arrest and myocardial infarction.

About this issue a participant said: "man's life depends on heart; if a problem is made, the whole life is endangered, one of our neighborhoods had heart problem for years, the poor man had no relaxation, I am really not calm now". Another participant said: "Those early days when I was told that I have a heart problem and should be hospitalized, I didn't believe it at all. I didn't have any problem. I was confused and believed that my life was ruined".

2. Trying to Achieve Calmness through Spirituality

Following to become aware about the critical situation after the diagnosis of cardiovascular disease and discharge from hospital, the participants had concluded that they should keep calm to continue life. They utilized many strategies to keep calm. Many of them had resorted to spirituality and tendency to religious practices to keep calm.

About this issue a patient said: "After it became clear that I have heart disease I was trying to have night prayer, I woke up and spoke to God for one or two hours. I was calming myself down with this plan. Through this I was keeping my morale high. I could remove my anxious by reading Quran and saying prayer".

About this issue another patient said: *“After I was discharged, I told O Lord! You created us, our deaths and lives are in your hands, ”I just put my trust in God.*

A patient said about his prayers after heart disease: *“I was only thinking of God, I prayed, I had confided myself to God and thanked God”.*

3. Losing Health

Many of participants said they have not paid much attention to their health status before diagnosis of the disease and before appearance of the signs and symptoms of cardiovascular diseases. They had never thought that they might have cardiovascular problems. Having had disclosure of the disease, they felt that they had lost their health and each moment were waiting for its aftermath. The patients mentioned the importance of heart and considered its health as health of the body. One of the patients said: *“with this heart disease I know I have lost my health, each moment my heart may become more clogged and have worse status. If man’s heart is clogged he will suffer from lesion”.*

In many participants' opinion, heart is the most sensitive organ of a body and needs high care and attention, but many of them didn't pay attention to the issue before suffering from heart disease at all.

Another patient said: *“Before suffering from the disease I didn't know about sensitivity and vulnerability of heart a lot, if I had suitable food diets and exercised, maybe I did not suffer from heart disease now and I was healthy and peppy”.*

4. Trying to Promote Health through Lifestyle Modification

After concluding that they need to promote their health, participants had used several strategies. They decided to modify their lifestyles to promote their health and hearts.

About this issue, one of the patients said: *“man should do something before a problem happens, now that I have this situation I should try hard to maintain my health”.*

Some participants mentioned instances on lifestyle modification: following suitable food and drug diets, regular referring to hospital in order to control cardiovascular status, managing stress, modifying work hours, regular and planned exercising, walking, and positive thinking.

A patient said about his decision to have further activity after diagnosis of heart disease: *“I was very busy and I also wasn't aware a lot, I did not know about the importance of the issue. For example, you should exercise an hour a week, control nutrients, and avoid cooking foods with high saturated oils. I used to*

go mountains once a week for one or two hours ,but this was not enough for a person who is sitting all the time, should deal with customers, becomes angry regularly, and needs to work hard”

Another patient said: *“After diagnosis of the disease, I decided to have activity and do exercise, I shouldn't sit in a place like old men, the situation was difficult”*

Another participant mentioned his daily exercises and said: *“After my heart disease, I exercised every day I woke up at 5a.m., went to park and exercised.”*

Another patient emphasized the necessity to take care of heart and said: *“After the disease, heart needs care and the suitable care is to avoid consuming salt, fat, and sweets according to physicians' orders. Doctors gave us an instruction which we should observe, such as not eating salt, fat, and red meat. I observed these cases”.*

5. Perceiving Family Support

Participants mentioned that they perceived family support more than before. They considered that family support from the patient is important and emphasized that this patient's life depends on support of family. In this regard a participant said: *“family support from a patient with heart disease is very important. If the family doesn't support the patient, he/ she become effete and will die”.*

A participant mentioned the effect of support from a patient with heart disease and said: *“After the disease my family supported me a lot. If there is no support, the patient will suffer a worse disease. When I see my family is sympathetic, supportive, as well as cooperative and I have some facilities, I become motivated”.*

6. Living in Fear

For patients living with coronary artery disease was always accompanied with fear, anxiety, uncertainty, fear of the future, and feelings of hopelessness so that all participants after suffering from cardiovascular diseases expressed such experiences. They were always afraid of heart attack, cardiac arrest, and sudden death. Being worry and having fear were so severe in some patients that they were worry about heart attack even in sleep. They were afraid of cardiac arrest during nights; feeling worry and fear caused disturbance in their sleeps.

In this regard, a patient said: *“After heart disease I had much concern, I was too sad. Sometimes I slept and said I will die tomorrow morning and will not wake up”.*

About living after heart disease a patient mentioned: *“living with heart disease is with anxiety and worry; being worry about cardiac arrest and*

death”

“My life after heart disease has always been accompanied with anxiety, I’m thinking if I sleep in night I will have rupture of heart tomorrow morning. I’m always worried, I’m always sad” said a patient about his living after heart disease.

Another participant talked about feeling of being hopeless after suffering from heart disease: *“After heart disease I was always assuming two modes for myself and the situation was a new experience for me. I was in a situation which my hope to continue living was very low”*.

A patient expressed his uncertainty about his heart: *“after the disease my heart is not working correctly, there is no confidence to that. When a heart becomes ill, you should take care of yourself a lot, you need to limit yourself.”*

7. Living with a Damaged Heart

Some participants said their hearts have been damaged after suffering from coronary artery disease and should live with damaged hearts for the rest of their lives. They mentioned some symptoms such as chest pain, shortness of breath, edema, palpitations, and cough.

A participant talked in this regard: *“After the disease my heart was damaged; now I’m living with a damaged heart.”*

Another participant said about his heart status after the disease: *“If a human’s heart is damaged what is left for him/her to live further? I didn’t care about myself and damaged my heart.”*

Discussion

Findings of the research showed that the lived experiences of elderly patients with coronary artery disease were with “ups and downs” with sub-themes of “losing the calmness, trying to achieve calmness through spirituality, losing health, trying to promote health through modification of lifestyle, perceiving family support, living in fear and living with a damaged heart”. Living with coronary artery disease was with ups and downs for participants of the study. They were frequently comparing living with coronary artery disease to their normal status. Suffering from the disease had divested patients from calmness and confidence; they were expressing some concerns about future by imagining probability of having heart problems in future such as heart attack and cardiac arrest was with fear and stress. Many studies had mentioned heart disease as a stressful and anxious event (5). Symptoms of cardiovascular diseases are with anxiety, depression, and fear of physical disability; this factor causes losing calmness of the person by sudden and unexpected start.

Participants were using many strategies to keep calm and return to the normal situation. They utilized the strategy of spirituality, tendency to religious actions, and paid more attention to spiritual issues to keep and promote calm. Positive effect of suffering from coronary artery disease in elderly participants was the fact that they utilized several strategies. Considering the cultural and religious grounds of patients and also their religions, i.e., all of them were Shia, they all used spirituality and religion to keep calm and achieve calmness. Spirituality and tendency to spirituality are also experienced in patients with cancer, cardiovascular diseases, and chronic disease (3-17). Momen Nasab et al., mentioned “trust in God” in their research on patients' experiences. Also, they added that heart attack makes spiritual dimension of participants more prominent and spirituality performs an important role to deal with such stressful and dangerous events of heart disease. They had expressed that spirituality had a significant effect on their recovery process from the starting moments of the disease towards recovery process (5).

All patients mentioned roles of their religious beliefs in bearing hardships and problems after coronary artery disease. Many researches have shown that patients use their religious beliefs in encountering problems and difficulties. Spiritual beliefs are to feel dependency on a divine and powerful force which causes life satisfaction and reduction of depression in patients (18, 19).

Participants were adapted to their problems by remembering power of God in all affairs of life and by resorting to Quran and Imams. They were always mentioning the holy verse of “Remembrance of God certainly brings comfort to all hearts” “Verily hearts become calm with remembrance of God”. Using spirituality in addition to compatibility with problems had effects on recovery process. In many studies the relation of spirituality, physical and psychological-spiritual health, as well as compatibility with disease has been mentioned (5, 14, 17).

After experiencing coronary artery disease, patients concluded that they have lost their health and had to try to achieve it. All their efforts were in order to reach health again and even promote it through some strategies. They had decided to modify their lives to promote their health. They were utilizing some strategies such as managing stress, regular exercising, modifying working hours, and changing food diet. After a heart event, patients had no other way to modify lifestyle.

Afrasiabi far et al., wrote: “elderly people had used some approaches to reduce consequences of the disease and return to previous life after heart attack despite tensions and limitations. They believed returning to normal life requires maintenance of abilities despite the existing limitations. For this purpose, patients had considered self-reliance and self-care as their strategies (20).

Patients said after suffering from coronary artery disease, they had well been supported by family. They were aware of the critical role of family support and thought that continuing their lives with the disease

mostly depends on family support. Some experts described family support as the level of having affection, companionship, care, respect, attention, and received help by other people, groups, friends, and others. Patients also mentioned that they received affection, companionship, support, and encouragements from their families.

Attention of family to patient is one of the most essential kind of care, according to Iranian culture, since families have deep relationships with members. Many reports are available about effects of families' supports on patients. Supporting patients prevents from the outbreak of physiological effects of disease, increases self-care, and has a positive effect on physical and social statuses of patients. Also, family support increases patient's performance (5, 21, 22).

For patients, living with coronary artery disease was always accompanied by fear, anxiety, worry so that all participants had experienced fear after disclosure of the disease. They were always afraid of heart attack, cardiac arrest, and sudden death. The prevalence of worry and fear of heart attack were so high in some patients that they felt them even in sleep. The negative feelings of anxiety and fear caused disturbance of their sleeps. Afrasiabi Far et al., expressed experience of patients' lives after heart attack as "in shadow of fear". Theme of life in shadow of fear had sub-classes of death fear by having another heart attack, fear of physical disability, and fear of becoming dependent to others. Fear of cardiac arrest and sudden death is one of psychological responses of elderly patients with heart attack which they experience from beginning of symptoms (20, 23). Experience of fear included fear of death during sleep and fear of pain during going out from house (3, 24). Participants mentioned importance of heart in lifetime. They mentioned heart as the "king of body". This feeling that heart is damaged now made them to be always worry. They suffered from fear, worry, and feeling that they should face death very soon due to damaged heart. In a research, "seeing death" and "the verge of death" were reported as heart disease themes (3, 25).

Patients had feelings of hopelessness about future of their lives following feelings of damaged heart and mental involvement. This experience was challenging for them and made them worry that their physical statuses may become worse every moment. Feelings of hopelessness had affected all aspects of their lives (8). Fear of disability to do the routine actions and dependency to others were most prevalent fears of most patients.

Conclusion

Findings of the research indicated lived experiences of elderly patients with coronary artery disease. The coronary artery disease had made them to lose their calmness. They were trying to achieve calmness by tending to spirituality and doing religious activities. Aware of coronary artery disease, patients imagined that they have lost their health and via modifying lifestyle, they were trying to promote their health.

They were supported by family along with their claim of high positive effects of family members on heart disease. Their lives was with fear, anxiety, worry, and feeling of hopeless. They imagined that their hearts are damaged and should continue their lives with damaged heart. All findings of this study emphasized on necessity of training patients and their family members. Also, findings of the research can be used in care, training, and managing fields. Therefore, professors and trainers of nursing, nurses working in clinic and society, nursing managers, patients' planners and families can use findings of the research to promote patients' training, consulting, quality of care and support from patients more and more.

It is suggested to conduct a study in the field of strategies of achieving calmness and effects of religious consultations on promoting calmness and giving confidence to patients' lives suffering from coronary artery disease.

Study limitations

Some limitations of qualitative studies are small sample size and the inability to generalize the results. The forgetfulness of some elderly participants about their experiences can be considered as another limitation of this research.

Conflict of interest

None.

Acknowledgment

This article is resulted from the approved projects of Shahed University IR. Shahed. REC 1395.108. Hereby, the financial support of the university and participants' cooperation in the study are appreciated.

References

1. American Diabetes Association. Cardiovascular disease and risk management. *Diabetes Care*. 2016; 39(Supplement 1): 60-71.
2. McAloon CJ, Boylan LM, Hamborg T, Stallard N, Osman F, Lim PB, et al. The changing face of cardiovascular disease 2000–2012: An analysis of the world health organisation global health estimates data. *International Journal of Cardiology*. 2016; 224: 256-64.
3. Abbasi M, Mohammadi N, Nikbakht Nasrabadi A, Sadegi T. Experiences of Living with Coronary Artery Bypass Graft: a Qualitative Study. *Journal of Hayat*. 2014; 19(4): 38-47. [Persian]
4. Heusch G, Libby P, Gersh B, Yellon D, Böhm M, Lopaschuk G, et al. Cardiovascular remodelling in coronary artery disease and heart failure. *The Lancet*. 2014; 383(9932): 1933-43.

5. Momennasab M, Moattari M, Abbaszade A, Shamshiri B. Spiritual experience of heart attack patients: A qualitative study. *Journal of Qualitative Research in Health Sciences*. 2013; 1(4): 284-97. [Persian]
6. Shafipour V, Mohammadi E, Ahmadi F. Experience of open heart surgery patients from admission to discharge: a qualitative study. *Iranian Journal of Critical Care Nursing*. 2013; 6(1): 1-10. [Persian]
7. O'Keefe-McCarthy S, McGillion MH, Victor JC, Jones J, McFetridge-Durdle J. Prodromal symptoms associated with acute coronary syndrome acute symptom presentation. *European Journal of Cardiovascular Nursing*. 2016; 15(3): 52-9.
8. Mohammadi N, Abbasi M, Nasrabadi AN, Salehiomran A, Davaran S, Norouzadeh R. Passion for life: lived experiences of patients after coronary artery bypass graft. *The Journal of Tehran University Heart Center*. 2015; 10(3): 129.
9. Arthur HM, Wright DM, Smith KM. Women and heart disease: the treatment may end but the suffering continues. *Canadian Journal of Nursing Research Archive*. 2016; 33(3): 17-29.
10. Azizi F, Rahmani M, Emami H, Mirmiran P, Hajipour R, Madjid M, et al. Cardiovascular risk factors in an Iranian urban population: Tehran lipid and glucose study (phase 1). *Sozial-und Präventivmedizin/Social and Preventive Medicine*. 2002; 47(6): 408-26.
11. Moeini M, Naseri N, Zargham-Boroujeni A. Lived experiences of women with ischemic heart disease. *Iranian Journal of Nursing and Midwifery Research*. 2012; 17(2):137-42.
12. Van Manen M, editor. *Writing in the dark: Phenomenological studies in interpretive inquiry*. Left Coast Press; 2003.
13. Taheri ZS, Zandi MA, Ebadi A. The effect of spiritual care based on Ghalbe Salim Model on the sleep quality of the patients with coronary artery disease. *Iranian Journal of Critical Care Nursing*. 2014; 7(2): 92-101.
14. Zamanzadeh V, Rassouli M, Abbaszadeh A, Nikanfar A-R, Alavi-Majid H, Mirza-Ahmadi F, et al. Spirituality in cancer care: a qualitative study. *Journal of Qualitative Research in Health Sciences*. 2014; 2(4): 366-78.
15. Akhbardeh M. Role of spiritual beliefs and prayer in health promotion of chronic patients: a qualitative study. *Quran and Medicine*. 2011; 1(1): 5-9.
16. Abbasi M, Tasooji HR, Hasan M, Ahmari Tehran H, Sadeghi T, Dehghani F, et al. Exploring the hospitalized patients' religious expectations of nurses; a qualitative study. *Health, Spirituality and Medical Ethics*. 2016; 3(3): 30-6.
17. Abbasi M, Mohammadi N, Nikbakht nasrabadi A, Gaeeni M. Towards spirituality after coronary artery bypass grafting: A hermeneutic phenomenological study. *Health, Spiritual and Medical Ethics*. 2017; 4(1): 10-15.
18. Cowlshaw S, Niele S, Teshuva K, Browning C, Kendig H. Older adults' spirituality and life satisfaction: a longitudinal test of social support and sense of coherence as mediating mechanisms. *Ageing and Society*. 2013; 33(07): 1243-62.
19. Nourisaeed A, Salari A, Nourisaeed A, Rouhi Balasi L, Moaddab F, Akbari B. Comparison of religious attitudes in patients with coronary artery disease and healthy people. *Iranian Journal of Nursing Research*. 2015; 10(3): 18-25. [Persian]
20. Falahi Khoshknab M, Hasani P, Afrasiabifar A, Yaghmaei F. Iranian elderly experiences with myocardial infarction. *Iranian Journal of Ageing*. 2008; 3(3): 66-74. [Persian]
21. Najafi Ghezalje T, Yadavar Nikraves M, Emami A. Coronary heart disease patients transitioning to a normal life: Perspectives and stages identified through a grounded theory approach. *Journal of Clinical Nursing*. 2014; 23(3-4): 571-85.
22. Lisa W, Amanda T, Ma'en Aq, Amanda Ch. The role of the family in supporting the self-management of chronic conditions: A qualitative systematic review. *Journal of Clinical Nursing*. 2017: 1-9.
23. Fors A, Dudas K, Ekman I. Life is lived forwards and understood backwards—experiences of being affected by acute coronary syndrome: a narrative analysis. *International Journal of Nursing Studies*. 2014; 51(3): 430-7.
24. Xavier D, Gupta R, Kamath D, Sigamani A, Devereaux P, George N, et al. Community health worker-based intervention for adherence to drugs and lifestyle change after acute coronary syndrome: a multicentre, open, randomised controlled trial. *The Lancet Diabetes & Endocrinology*. 2016; 4(3): 244-53.
25. Hildingh C, Fridlund B, Lidell E. Access to the world after myocardial infarction: experiences of the recovery process. *Rehabilitation Nursing*. 2006; 31(2): 63-8.