



## Original Article

# Association between Chronic Pain and Depression among the Elderly of Amirkola City, Northern Iran

Alijan Ahmadi Ahangar<sup>1\*</sup>, Seyed Reza Hosseini<sup>2</sup>, Farzan Kheirkhah<sup>2</sup>, Masoumeh Karimi<sup>3</sup>, Payam Saadat<sup>4</sup>, Ali Bijani<sup>2</sup>, Shayan Alijanpour<sup>5</sup>, Hamed Hosseinzadeh<sup>1</sup>, Sakineh Faraji<sup>1</sup>

<sup>1</sup>. Mobility Impairment Research Center, Babol University of Medical Sciences, Babol, Iran

<sup>2</sup>. Social Determinants of Health Research Centre, Babol University of Medical Sciences, Babol, Iran

<sup>3</sup>. Student Research Committee, Babol University of Medical Sciences, Babol, Iran

<sup>4</sup>. Ayatollah Rouhani Hospital, Babol University of Medical Sciences, Babol, Iran

<sup>5</sup>. Emergency Medical Service Center, Babol University of Medical Sciences, Babol, Iran

## ABSTRACT

### Article history

Received 13 Jul 2017

Accepted 19 Sep 2017

**Introduction:** Chronic pain and depression are two of common geriatric disorders. The aim of this study was to investigate the association between chronic pain and depressive symptoms in older people in Amirkola.

**Methods:** This cross-sectional study is part of Amirkola Health and Aging Project. It is an ongoing cohort project which was being conducted from 2011 on all people aged 60 and above in Amirkola city in northern part of Iran. One thousand six hundred and four older people were included in the study. Data related to chronic pains and their locations were collected using questionnaire by inquiring the elderly while depressive symptoms were gathered based on Geriatric Depression Scale. Data were analyzed by chi-square test and logistic regression in SPSS.

**Results:** The prevalence of chronic pain, depressive symptoms, and co-occurrence of chronic pain and depressive symptoms among the elderly was respectively 84.4%, 43.5%, and 39.8%. The odds ratio of having chronic pain in depressed people was more than non-depressed people (OR = 2.88; 95% CI = 2.11-3.94). Prevalence of chronic pain in hands, wrists, elbows, shoulders, neck, hip joints, knees, ankles, legs, and back were noticeable in people with symptoms of depression compared to those without depressive symptoms. In regression model, severity of depressive symptoms (OR = 1.73 (1.23 - 2.45)), being female (OR = 2.40 (1.68 - 3.45)), increasing age (OR = 1.03 (1.01 - 1.05)) and having chronic diseases (OR = 1.24 (1.13-1.35)) were among the important variables that influenced chronic pain.

**Conclusion:** This study showed, especially in women, significant association between chronic pain and depressive symptoms. It is essential to take prophylactic and treatment measures suitable for their control and treatment.

**Keywords:** Depression, Chronic Pain, Aged

**Citation:** Ahmadi Ahangar A, Hosseini SR, Kheirkhah F, Karimi M, Saadat P, Bijani A, et al. Association between chronic pain and depression among the elderly of Amirkola city, northern Iran. *Elderly Health Journal*. 2017; 3(2): 74-79.

### Introduction

Increased life expectancy has been one of the great achievements of the new century. About 700 million people of the world's population are the elders above 60 years of age and by 2020, this figure will amount to more than one billion (1). According to the

Statistical Center of Iran, it is expected that the number of elders in Iran reaches more than 25 million by 2050 (2). Chronic pain is a major problem in public health. They affect physical and mental health of people and reduce their quality of life (3).

\* **Corresponding Author:** Mobility Impairment Research Center, Babol University of Medical Sciences, Babol, Iran. Tel: +989111111202, Email address: ahmadiyahangaralijan@yahoo.com

The International Association for the Study of Pain has defined chronic pain as a pain with no apparent biological value and without severe tissue lesions that extends beyond the time of the treatment of a disease. Generally, pains lasting more than 3 months are also classified as chronic pain (4). The most common chronic pain in the elderly include musculoskeletal pain (joints, the lower and the lower extremities). Studies also show that the most important mental illness among the elderly is depressive disorder (6). Depression is known as one of the most common sensational and emotional problems among elders (7). There is much controversy over the relationship between depression and chronic pain. Some studies support the idea that depression puts people at the risk of developing chronic pains. On the other hand, depressed patients report significant degrees of pain (8). The prevalence of chronic pain in patients who are under treatment for depression is reported between 51.8 to 59.1% (9). On the other hand, some research supports the theory that depression appears following chronic pains (10). Some other studies assume a mutual relationship between pain and depression. This is because depression and chronic pain interact with each other in a way that worsens the other (11). Some researchers believe that the presence of both pain and depression may be assumed as a pain-depression syndrome rather than the existence of two independent disorders (12). However, depression imposes high burden on patients with chronic pain, their family and care giver, and health systems. In chronic pains, depression increases the costs related to disability and use of health care (13). Having considered that depression, in addition to being one of the diseases associated with chronic pain at least in 13% of the elderly, it also is one of risk factors of chronic pain in elderly (4). The main objective of this study was to assess the association of chronic pains and depression among the elderly of Amirkola City.

## Methods

### *Participants and the study protocol*

This cross-sectional descriptive study came from the Amirkola Health and Aging Project (AHAP) which was being conducted from 2011 on all people aged 60 and above in Amirkola city (14). From the total of 2,234 people aged 60 and above, 1616 people participated in this study (response rate = 72.3%). By providing information on the study, people were invited to participate in the study through telephone calls and home visits. With considering exclusion criteria, 1604 older adults consisting of 876 men and 728 women were included in the present study.

All people aged 60 and above in Amirkola who completed all questionnaires regarding study variables were entered in the study. Moreover those who had mental illnesses or cases such as cognitive impairments were excluded.

### *Measures*

The data were collected using a questionnaire composed of demographic characteristics including: age, gender, drugs, and history of various diseases. Chronic musculoskeletal pain and its site were assessed using a questionnaire. The participants were asked whether they had experienced persistent pain (lasting for at least three months) within the past six months (15) in the following sites: hands, wrists, elbows, shoulders, face, jaw, neck, hips, knees, ankles, legs and back. All questionnaires were completed through conducting interviews with the participants. The presence of depressive symptoms in the elderly was examined using a standard Geriatric Depression Scale (GDS) questionnaire validated previously in Iranian population which contains 15 questions and each question rates one point (16-18). Ten questions indicate the presence of depression when answered positively, while the rest (question numbers 1, 5, 7, 11, 13) indicate depression when answered negatively. A cut-off of five or more symptoms was defined as having depressive symptoms. Depressive symptoms were divided into mild (5-8), moderate (9-12), and severe (13-15). Using this cut-off, the test sensitivity was 92% and specificity was 89% (4, 16). The Cronbach's alpha of this questionnaire was 0.81 in the elderly population of Amirkola.

To evaluate cognitive status, we used Mini Mental State Examination. It contains 11 items and evaluates five cognitive functions, including orientation, attention, memory, language and visual-spatial skills. The maximum score is 30 points and individuals with score of 25 or lower indicate the presence of cognitive impairment (19). The validity and reliability of this test for Iranian patients have already been approved (20).

Serum vitamin D levels (25-hydroxy vitamin D) were measured through laboratory evaluation of morning blood samples of the elderly using ELISA test in Cellular and Molecular Biology Research Center, Babol University of Medical Sciences.

### *Ethical considerations*

This study was approved by the Ethics Committee of the Research Institute of Babol University of Medical Sciences (ethic number: 2707). All the elders who participate in AHAP were taken informed consent.

### *Data analysis*

Collected information was entered into SPSS and analyzed by chi-square test to determined possible association between the cases in question. Data were also analyzed using logistic regression to determine the association of chronic pain in different areas with symptoms of depression. P-Value <0.05 was considered significant.

## Results

Of 1604 older adults were studied, 876 males (54.6%) and 728 were females (45.3%). The mean age of respondents was  $69.35 \pm 7.41$  years (the range of 60 to 92), and the majority of people were in the age group of 60 to 64 years. The prevalence of chronic pain among the elderly of Amirkola was 84%. The prevalence of chronic pain by sex was statistically significant (684 (78.1 %) male vs. 670 (92 %) female,  $p < 0.001$ ). The prevalence of depressive symptoms among the elderly was 43% and significant difference with sex (256 (29.2 %) male vs. 441(60.6 %) female,  $p < 0.001$ ). Moreover, associated chronic pain with depressive symptoms among the elderly of Amirkola was 638 (39.8%) and difference was significant by sex (222 (25.3%) male vs. 416 (57.1%) female ( $p = 0.000$ )). The risk of chronic pain in older adults with depressive symptoms, both in general and in each of the two genders, was significantly higher than non-depressed elders. In general, the risk of chronic pain among depressed people was 2.88 times higher than non-depressed ones (CI = 2.11-3.93,  $p < 0.001$ ). This risk was 2.16 (CI = 1.25 - 3.72,  $p = 0.005$ ) in female and 2.23 (CI = 1.49 - 3.34,  $p < 0.001$ ) in men. The Results showed that with increase in depression scale, prevalence of chronic pain increases in total and female (Table 1). There was a significant association between the prevalence of chronic pain in different parts of the body and symptoms of depression (except in chronic pain where the information was insufficient). The risk of chronic pain in any of these areas among depressed people was higher than non-depressed people. For example, the risk of chronic pain in hand among depressed patients was 2.34 times higher than those without depressive symptoms (CI = 1.85 - 2.96,  $p < 0.001$ ) (Table 2).

The logistic regression analysis confirmed that having depressive symptoms, being female, age and number of chronic diseases were risk factors and vitamin D had protective effect for chronic pain (Table 3).

## Discussion

The prevalence of chronic pain among the elderly of Amirkola was 84.4 % that statistically significant in female while this trend was less significant in European studies (31.8% to 69.8%) (21-22). This sexual disproportionality can be attributed to sex hormones, endogenous pain modulation system, and cognitive/emotional factors between male and female.

In this study, the chronic pain was associated with depression in 39.8% cases which was statistically more prominent in females than males (57.1% versus 25.3%). Also in this study, with increasing in severity of depressive symptoms, prevalence of chronic pain increased that was statistically significant in females. In other studies, the prevalence of chronic pain increased in accordance with increasing depression scale (21-22).

Differences between female and male regarding the prevalence of chronic pain and associated depression obtained in this study were consistent with Calvo et al. study (22). Females may express symptoms more than males. This causes psychological distress and raises the risk of chronic pain and associated depression (22). Chronic knee pain (63%), back pain (45.4%), and pain in legs (43.5%) had the highest prevalence, because of most people are farmer that is considered as a heavy activity in this area.

Furthermore, the results indicate a significant association between depression and chronic pain in the studied areas (except jaw and face), and the risk of chronic pain in these areas among people with depression symptoms was higher than non-depressed people. As the results show, 71.3% of people with depressive symptoms had chronic knee pain such that the risk of chronic knee pain in people with depressive symptoms was 1.91 times higher than individuals without symptoms of depression that consistent with Han study (23).

**Table 1. Distribution of chronic pain in the Amirkola elderly, according to subgroups of GDS score by gender**

GDS Score		Without chronic pain N (%)	With chronic pain N (%)	P
<b>Total</b>	<b>Normal</b>	191 (21.1)	716 (78.9)	< 0.001
	<b>Mild depression</b>	40 (9.1)	398 (90.9)	
	<b>Moderate depression</b>	14 (7.9)	163 (92.1)	
	<b>Severe depression</b>	5 (6.1)	77 (93.95)	
<b>Female</b>	<b>Normal</b>	33 (11.5)	254 (88.5)	0.021
	<b>Mild depression</b>	18 (7.1)	237 (92.9)	
	<b>Moderate depression</b>	5 (4.15)	116 (95.9)	
	<b>Severe depression</b>	2 (3.1)	63 (96.9)	
<b>Male</b>	<b>Normal</b>	158 (25.5)	462 (74.5)	< 0.001
	<b>Mild depression</b>	22 (12)	161 (88)	
	<b>Moderate depression</b>	9 (16.1)	47 (73.9)	
	<b>Severe depression</b>	3 (17.6)	14 (82.4)	

**Table 2. The risk of chronic pain in different parts of the body in terms of depressive symptoms among the elderly of Amirkola, Northern Iran**

Region	Subgroup	Individuals with chronic pain N (%)	Individuals without chronic pain N (%)	Odds ratio	95%CI	P
Hand	Depressed	226 (32.4)	471 (67.6)	2.34	(1.85-2.96)	< 0.001
	Non-depressed	154 (17)	753 (83)	1	-	
Wrist	Depressed	81 (11.6)	616 (88.4)	2.16	(1.50 -3.10)	< 0.001
	Non-depressed	52 (5.7)	855 (94.3)	1	-	
Elbow	Depressed	105 (15.1)	592 (84.9)	3.65	(2.51-5.30)	< 0.001
	Non-depressed	42 (4.6)	865 (95.4)	1	-	
Shoulder	Depressed	320 (45.9)	377 (54.1)	1.87	(1.52 -2.29)	< 0.001
	Non-depressed	283 (31.2)	624 (68.8)	1	-	
Face	Depressed	5 (0.7)	692 (99.3)	1	(1.01-1.02)	0.011
	Non-depressed	0 (0)	907 (100)	1	-	
Neck	Depressed	222 (31.9)	475 (68.1)	1	(1.65-2.63)	<0.001
	Non-depressed	166 (18.3)	741 (81.7)	-	-	
Hip joints	Depressed	215 (30.8)	482 (69.2)	2.08	(1.64 -2.63)	< 0.001
	Non-depressed	160 (17.6)	747 (82.4)	1	-	
Knees	Depressed	497 (71.3)	200 (28.7)	1.90	(1.54 -2.35)	< 0.001
	Non-depressed	513 (56.6)	394 (43.4)	1	-	
ankles	Depressed	90 (12.9)	607 (87.1)	1.82	(2.54-1.31)	< 0.001
	Non-depressed	68 (7.5)	839 (92.5)	1	-	
Legs	Depressed	357 (51.2)	340 (48.8)	1.74	(1.42-2.12)	< 0.001
	Non-depressed	341 (37.6)	566 (62.4)	1	-	
Back	Depressed	368 (52.8)	329 (47.2)	1.69	(2.06-1.38)	< 0.001
	Non-depressed	361 (39.8)	546 (60.2)	1	-	

**Table 3. Odds ratio and 95% confidence interval for associations between depressive symptoms and chronic pain in older people in Amirkola**

variables	OR	95%CI	P
Depressive symptoms	1.73	1.23-2.45	0.002
Gender	2.40	1.68-3.45	< 0.001
Age	1.02	1.01-1.04	0.026
Cognitive impairment	1.02	0.67-1.07	0.312
Social support	0.95	0.90-1.01	0.061
Vitamin D level	0.49	0.26-0.90	0.023
Number of chronic diseases	1.23	1.12-1.35	< 0.001

Phyomaung study showed a strong association between depression and knee pain and indicated that depression plays a major role in knee pain (24). In Kroenke study change in depression severity was diagnosed as a strong predictor of knee pain severity following that (25). Moreover, 52.8% of people with symptoms of depression suffered chronic back pain. Hence, the risk of chronic back pain in people with depressive symptoms versus those without such symptoms was 1.692. The results are consistent with other studies in this field. Glombiewski study showed higher degrees of depression were associated with more severity of pain and pain-related disability and more than one third (36.4%) of them had depression and pain comorbidity (26). In Stubbs study there was a significant association between severity of depression and back pain (OR = 2.21) (27). Pinheiro study showed that depressive symptoms increase the risk of developing back pain (OR = 1.59; 95% CI = 1.26-2.01) (9, 28-29). In Gerrits study there was a significant difference between pain in the face, mouth and depression (11).

### Conclusion

In conclusion, there was a strong association between chronic pain and depression. In addition, the prevalence of chronic pain, depression, and chronic pain and depression in female was significantly higher than men. Given the high prevalence of depression, pain and comorbidity among the elderly of Amirkola compared to the international community, it is necessary to take measures and treatment procedures to control and treat them.

### Study limitations

This study has a few limitations that should be mentioned. The cross-sectional nature of the study that makes difficult to find causal relationships. Data about variables were not based on clinical findings and self-reported data might lower its accuracy.

### Conflict of interest

There was no conflict of interest.

### Acknowledgement

We hereby appreciate the Deputy for Research and Technology of Babol University of Medical Sciences for approval and financial support of the mentioned proposal. We also thank the officials of the Master Plan of "Amirkola Aging and Health Project" (AHAP) to the number 892917 and all the elderly who appreciated in this study.

### Authors' contribution

Study design: AAA, SRH, FK, PS

Interpretation of data: AAA, SRH, PS, AB, SA

Acquisition of data: SRH, FK, MK, SA, HH, SF

Revising the article: AAA, SRH, FK, MK, SA, HH, SF

Analysis: AB

Final approval: AAA, SRH

### References

- Ghaneh B, Saeed-Banadaky SH, Rahaei Z, Rezaeipandari H, MohitiArdakani E. Disability and Self-Care among Elders in Yazd. *Elderly Health Journal*. 2016; 2(1): 39-44.
- Shahbazi MR, Mirkhani M, Hatamizadeh N, Rahgozar M. Evaluation of disability in Tehran elderly. *Iranian Journal of Ageing*. 2009; 3(3-4): 84-92. [Persian]
- Gaskin DJ, Richard P. The economic costs of pain in the United States. *The Journal of Pain*. 2012; 13(8): 715-24.
- Zis P, Daskalaki A, Bountouni I, Sykioti P, Varrassi G, Paladini A. Depression and chronic pain in the elderly: links and management challenges. *Clinical Interventions in Ageing*. 2017; 12: 709-20.
- Makris UE, Abrams RC, Gurland B, Reid MC. Management of Persistent Pain in the Older Patient A Clinical Review. *Journal of the American Medical Association*. 2014; 312(8): 825-36.
- Fiske A, Wetherell JL, Gatz M. Depression in Older Adults. *Annual Review of Clinical Psychology*. 2009; 5: 363-89.
- Duberstein PR, Pálsson SP, Waern M, Skoog I. Personality and risk for depression in a birth cohort of 70-year-olds followed for 15 years. *Psychological Medicine*. 2008; 38(5): 663-71.
- Demyttenaere K, Bruffaerts R, Lee S, Posada-Villa J, Kovess V, Angermeyer MC, et.al. Mental disorders among persons with chronic back or neck pain: results from the World Mental Health Surveys. *Pain*. 2007; 129(3): 332-42
- Gerrits MM, van Oppen P, van Marwijk HW, Penninx BW, van der Horst HE. Pain and the onset of depressive and anxiety disorders. *Pain*. 2014; 155(1): 53-9.
- Walker A.K, Kavelaars A, Heijnen C.J, Dantzer R. Neuroinflammation and Comorbidity of Pain and Depression. *Pharmacological Reviews*. 2014; 66(1): 80-101.
- Doan L, Manders T, Wang J. Neuroplasticity underlying the comorbidity of pain and depression. *Neural Plasticity*. 2015; 2015: 1-16.
- Li JX. Pain and depression comorbidity: a preclinical perspective. *Behavioural Brain Research*. 2015; 276(1): 92-8.
- Katon W. The impact of depression on workplace functioning and disability costs. *The American Journal of Managed Care*. 2009; 15(11): 322-7.
- Hosseini SR, Cumming RG, Kheirkhah F, Nooreddini H, Baiani MA, Mikaniki E, et al. Cohort.

- profile: The Amirkola Health and Aging Project. *International Journal of Epidemiology*. 2014; 43(5): 1393-1400
15. Treede RD, Rief W, Barke A, Aziz Q, Bennett ML, Benoliel R, et al. A classification of chronic pain for ICD-11. *Pain*. 2015; 156(6): 1003-7.
  16. Sheikh JI, Yesavage JA. Geriatric depression scale (GDS): Recent evidence and development of a shorter version. *Clinical Gerontologist: The Journal of Aging and Mental Health*. 1986; 5(1-2): 165-73.
  17. Malakouti SK, Fatollahi P, Mirabzadeh A, Salavati M, Zandi T. Reliability, validity and factor structure of the GDS-15 in Iranian elderly. *International Journal of Geriatric Psychiatry*. 2006; 21(6): 588-93
  18. Majdi M R, Ghayour Mobarhan M, Salek M, Taghi M, Mokhber N. Prevalence of Depression in an Elderly Population: A Population-Based Study in Iran. *Iranian Journal of Psychiatry and Behavioral Sciences*. 2011; 5(1): 17-24
  19. Folstein MF, Folstein SE, McHugh PR. "Mini-Mental State". A practical method for grading the cognitive state of patients for the clinician. *Journal of Psychiatric Research*. 1975; 12(3): 189-98.
  20. Seyyedian M, Fallah M, Nourozian M, Nejat S, Delavar A, Ghasemzadeh H. Validity of the Farsi version of mini-mental state examination. *Medical Journal of the Islamic Republic of Iran*. 2008; 25(4): 408-14.[Persian]
  21. Bauer H, Emeny RT, Baumert J, Ladwig KH. Resilience moderates the association between chronic pain and depressive symptoms in the elderly. *European Journal of Pain*. 2016; 20(8): 1253-65.
  22. Calvo-Perxas L, Vilalta-Franch J, Turró-Garriga O, López-Pousa S, Garre-Olmo J. Gender differences in depression and pain: a two year follow-up study of the survey of health, ageing and retirement in Europe. *Journal of Affective Disorders*. 2016; 193: 157-64.
  23. Han HS, Lee JY, Kang SB, Chang CB. The relationship between the presence of depressive symptoms and the severity of self-reported knee pain in the middle aged and elderly. *Knee surgery, sports traumatology, arthroscopy: official journal of the ESSKA*. 2015, 24(5):1634-1642
  24. PhyoMaung PP, Dubowitz J, Cicuttini FM, Fernando S, Wluka AE, Raaijmakers P, Wang Y, Urquhart DM. Are depression, anxiety and poor mental health risk factors for knee pain? A systematic review. *BMC Musculoskeletal Disorders*. 2014 ; 15: 10.
  25. Kroenke K, Wu J, Bair MJ, Krebs EE, Damush TM, Tu W. Reciprocal relationship between pain and depression: a 12-month longitudinal analysis in primary care. *The Journal of Pain*. 2011; 12(9): 964-73.
  26. Glombiewski JA, Hartwich-Tersek J, Rief W. Depression in chronic back pain patients: prediction of pain intensity and pain disability in cognitive-behavioral treatment. *Psychosomatics*. 2010; 51(2): 130-6.
  27. Stubbs B, Koyanagi A, Thompson T, Veronese N, Carvalho AF, Solomi M, et al. The epidemiology of back pain and its relationship with depression, psychosis, anxiety, sleep disturbances, and stress sensitivity: Data from 43 low-and middle-income countries. *General hospital psychiatry*. 2016; 43: 63-70.
  28. Pinheiro MB, Ferreira ML, Refshauge K, Ordoñana JR, Machado GC, Prado LR, et.al. Symptoms of Depression and Risk of New Episodes of Low Back Pain: A Systematic Review and Meta-Analysis. *Arthritis Care & Research*. 2015; 67(11): 1591-603.
  29. Tatebe M, Iwatsuki K, Hirata H, Oguchi T, Tanaka K, Urata S. Effects of depression and inflammatory factors on chronic conditions of the wrist. *Bone Joint Journal*. 2016; 98(7): 961-8.